

A rival for Babcock’s star: the extreme 30-kG variable magnetic field in the Ap star HD 75049*

V.G. Elkin¹, G. Mathys², D.W. Kurtz¹, S. Hubrig³, L.M. Freyhammer¹

¹*Jeremiah Horrocks Institute of Astrophysics, University of Central Lancashire, Preston PR1 2HE, UK*

²*European Southern Observatory, Casilla 19001, Santiago 19, Chile*

³*Astrophysical Institute Potsdam, D-14482, Potsdam, Germany*

19 October 2009

ABSTRACT

The extraordinary magnetic Ap star HD 75049 has been studied with data obtained with the ESO VLT and 2.2-m telescopes. Direct measurements reveal that the magnetic field modulus at maximum reaches 30 kG. The star shows photometric, spectral and magnetic variability with a rotation period of 4.049 d. Variations of the mean longitudinal magnetic field can be described to first order by a centred dipole model with an inclination $i = 25^\circ$, an obliquity $\beta = 60^\circ$ and a polar field $B_p = 42$ kG. The combination of the longitudinal and surface magnetic field measurements imply a radius of $R = 1.7 R_\odot$, suggesting the star is close to the zero-age main sequence. HD 75049 displays moderate overabundances of Si, Ti, Cr, Fe and large overabundances of rare earth elements. This star has the second strongest magnetic field of any main sequence star after Babcock’s star, HD 215441, which it rivals.

Key words: Stars: magnetic – stars: variables – stars: individual (HD 75049).

1 INTRODUCTION

The complex processes of atomic diffusion – radiative levitation and gravitational settling – occur in upper main sequence stars, and are particularly evident in chemically peculiar A, B and F-type stars, collectively known as Ap stars. Accordingly, these stars are good targets to place observational constraints on atomic diffusion, which is crucial for deepening our understanding of these processes within the framework of studies of the structure and evolution of stars, and for more global analysis of galactic chemical evolution. The Ap stars rotate considerably more slowly than spectroscopically normal main sequence stars with similar effective temperatures. A fraction of Ap stars have extremely long rotation periods from tens of days to tens of years, and even of the order of a century in the case of γ Equ. There is no clear picture of the braking mechanism leading to such slow rotation, but the observational evidence strongly suggests that it is related to magnetic field (Mathys et al. 1997; Abt 2009).

Many Ap stars possess strong, predominantly dipolar magnetic fields. The first magnetic star, 78 Vir, was found by Babcock (1947). Eleven years later Babcock (1958) pub-

lished the first catalogue of magnetic stars, and two years following that he discovered a huge 34 kG magnetic field in the star HD 215441 (Babcock 1960), known since then as “Babcock’s star”. Despite success in the years that followed in increasing the number known magnetic stars – especially in last several years (for example Hubrig et al. 2006; Kudryavtsev et al. 2006) – HD 215441 remains the record-holder for the main sequence star with the strongest magnetic field. The second strongest magnetic star, HD 154708, was discovered by Hubrig et al. (2005). Around the same time Kochukhov (2006) found HD 137509 to have magnetic field modulus of 29 kG. His result was in agreement with Mathys (1995) who discovered a very strong quadratic magnetic field up to 37 kG in this star and mentioned that it has the one of the strongest magnetic fields. Surprisingly, the longitudinal magnetic field in HD 137509 is not very large and varies just from -1.25 to $+2.35$ kG (Mathys 1991). HD 154708 still has the strongest magnetic field known among the cooler rapidly oscillating (roAp) stars (Kurtz et al. 2006).

The origin of the magnetic fields in Ap stars and details of their peculiar properties are still mysteries, despite large efforts in the study of these stars. As usual in physics, analysis of extreme cases may provide key information, hence there is great interest in the stars with the strongest magnetic fields to help solve many problems in Ap stars. Recently, Freyhammer et al. (2008) discovered an extremely

* Based on observations collected at the European Southern Observatory, Chile, in the programme 080.D-0170(A) and as part of programmes 078.D-0080(A) and 078.D-0192(A).

strong magnetic field in the Ap star HD 75049. Many lines in the high resolution spectra are split into clear Zeeman patterns. From the value of splitting Freyhammer et al. (2008) estimated the photospheric magnetic field modulus to be about 30 kG. These observations also revealed a reasonably short rotation period, and large variability both of the magnetic field and the spectral line intensities. Because of these interesting characteristics of this rare object, we have obtained a series of high resolution spectra and carried out circular spectropolarimetric observations with the VLT. We present in this paper the results of our analysis.

2 OBSERVATIONS AND DATA REDUCTION

To characterise the magnetic field of an Ap star, it is necessary to have magnetic measurements that cover the rotational phases of the star well. While we had a preliminary estimate for the rotation period of 4.05 d determined by Freyhammer et al. (2008) from All Sky Automatic Survey (ASAS) photometric survey data (Pojmanski 2002), we were uncertain about this, so decided to spread new observations over half a year to be sure to determine the correct rotational period. There is also a possible 5.28-yr period in the ASAS data that was noted by Freyhammer et al. (2008). The nature of this long period is not clear, but it is certainly not a rotational period; the $v \sin i = 8.5 \text{ km s}^{-1}$ of HD 75049 does not allow that. Further understanding of this period will require long-term observations.

A series high resolution spectra were obtained in service mode using UVES (Ultraviolet and Visual Echelle Spectrograph) on the ESO VLT (Very Large Telescope). Twelve spectra were collected between 2007 October 10 and 2008 April 1. Two previous observations of this star with FEROS and UVES obtained in 2007 February and March were also used for this analysis. These observations were made with echelle spectrographs over long spectral regions. For FEROS the spectral range is 3500–9220 Å with a resolution $R = \lambda/\Delta\lambda = 48\,000$; for UVES the range is 4970–7010 Å with a resolution of $R = 110\,000$. A journal of these observations is presented in Table 1. Software packages provided by ESO (UVES and FEROS pipelines in the MIDAS environment) were used for the reduction and extraction of 1D spectra.

We also obtained spectropolarimetric observations of HD 75049 with the **FO**cal **R**educer and low dispersion **S**pectrograph (FORS1) in the VLT in service mode during the time span 2007 December 30 to 2008 April 1. Observations with FORS1 spectra in circular polarisation are used for measurement of the longitudinal magnetic field, which corresponds to the line-of-sight component of the magnetic field vector integrated over the visible stellar hemisphere. This magnetic field parameter is sensitive to the magnetic field geometry and is very efficient for determining the rotational period.

Table 2 gives the Barycentric Julian Date of the middle of each observation, the exposure time, phase of the rotational period and the results of the longitudinal magnetic field measurements for three different lists of spectral lines or spectral regions. A detailed description of similar spectropolarimetric observations and its reductions is given in the recent paper by Hubrig et al. (2009).

Table 1. A journal of high resolution spectroscopic observations with UVES at the VLT and FEROS at the ESO 2.2-m telescope. The columns give the Barycentric Julian Date (BJD) of the middle of each exposure, the exposure time, the signal-to-noise ratio that was measured in sections of the continuum free of spectral lines.

BJD	exposure time sec	S/N ratio	Instrument
2454141.66963	1500	190	FEROS
2454171.50422	900	180	UVES
2454387.81593	1800	250	UVES
2454432.82489	1800	240	UVES
2454451.83366	1800	250	UVES
2454455.70226	1800	260	UVES
2454468.84589	1800	230	UVES
2454481.81465	1800	230	UVES
2454513.78892	1800	80	UVES
2454516.68904	1800	240	UVES
2454524.66461	1800	240	UVES
2454544.51608	1800	260	UVES
2454547.51127	1800	300	UVES
2454557.59095	1800	240	UVES

3 ROTATION PERIOD

The rotation period is an important parameter for the analysis of magnetic stars, which normally show photometric, spectral and magnetic variability with this period. In HD 75049 we found a clear picture of variability in all three cases. As mentioned above, photometric variability was detected in ASAS data, and spectral and magnetic variability were found from high resolution spectra (Freyhammer et al. 2008). Our new longitudinal field measurements have allowed us to determine the rotation period with confidence and higher accuracy.

Using a Discrete Fourier Transform (Kurtz 1985) and the programme PERIOD04 (Lenz & Breger 2005), we found a highest peak in the amplitude spectrum of the FORS1 B_z data (comprising 13 measurements) of $f = 0.2477 \text{ d}^{-1}$; using least squares fitting of a sinusoid we found $f = 0.2470 \text{ d}^{-1}$, as is seen in Fig. 1. We refined the fitted frequency and calculated uncertainties using nonlinear least-squares fitting, fitting the function

$$B_z = A_0 + A \cos[2\pi f(t - t_0) + \phi] \quad (1)$$

which gives our final values:

$$f = 0.246975 \pm 0.000005 \text{ d}^{-1}$$

$$P_{\text{rot}} = 4.04899 \pm 0.00008 \text{ d}$$

$$A_0 = -5127 \pm 34 \text{ G}$$

$$A = 4824 \pm 40 \text{ G}$$

$$\phi = 0.000 \pm 0.010 \text{ rad, where}$$

$$t_0 = \text{BJD } 245\,4509.550 \pm 0.006$$

$$\sigma = 121 \text{ G}$$

The standard deviation σ is per measurement with respect to the fit. The error on A_0 is thus $\frac{\sigma}{\sqrt{13}}$, where $N = 13$ data points. The error on t_0 is the error in phase divided by $2\pi f$. We thus get an ephemeris of

$$B_z^{\text{max}} = \text{BJD } 245\,4509.550 \pm 0.006 + 4.04899 \pm 0.00008 E. \quad (2)$$

The value of the rotation period is consistent with that de-

Table 2. Observations of HD 75049 with FORS 1. The columns give the Barycentric Julian Date (BJD) of the middle of each exposure, the rotational phase according to the ephemeris in eq. 2, and the longitudinal magnetic field for two wavelength ranges of the spectrum and for the hydrogen lines with the error of measurement.

BJD	rotation phase	longitudinal magnetic field B_z (G)		
		3212 – 6215 Å	3705 – 6215 Å	hydrogen lines
2454482.76787	0.3978	-8894 ± 68	-8934 ± 68	-9744 ± 132
2454483.60947	0.6057	-9071 ± 39	-9096 ± 39	-9938 ± 74
2454493.70464	0.0989	-905 ± 32	-902 ± 32	-1475 ± 61
2454516.73363	0.7865	-4471 ± 35	-4485 ± 35	-5064 ± 65
2454526.56586	0.2148	-3538 ± 40	-3564 ± 40	-3970 ± 74
2454527.58159	0.4657	-9832 ± 36	-9857 ± 36	-10529 ± 72
2454532.72343	0.7356	-5811 ± 49	-5824 ± 49	-6367 ± 91
2454539.77171	0.4764	-9639 ± 46	-9661 ± 46	-10334 ± 94
2454543.57545	0.4158	-9237 ± 39	-9257 ± 39	-9941 ± 75
2454546.52195	0.1435	-1821 ± 44	-1818 ± 44	-2364 ± 86
2454555.53106	0.3685	-8139 ± 33	-8168 ± 33	-8528 ± 64
2454557.61944	0.8843	-1785 ± 34	-1793 ± 34	-2397 ± 62
2454464.85968	0.9749	-640 ± 36	-639 ± 36	-1415 ± 68

terminated by (Freyhammer et al. 2008) from ASAS photometry.

4 THE STELLAR PARAMETERS

As an initial step to determine the fundamental stellar parameters we employed previous photometric observations. We used the *uvby* β photometry by Martinez (1993) and the UVBYBETA program written by T.T. Moon and modified by Napiwotzki et al. (1993) based on the grid published in Moon & Dworetsky (1985). We derived an effective temperature, $T_{\text{eff}} = 9600$ K, and a gravity of $\log g = 4.47$ (cgs). The surface gravity is an issue, since it is known that photometric calibrations of the c_1 index, which are useful for determination of luminosity (hence surface gravity) for normal stars, are not entirely suitable for Ap stars, where they tend to underestimate luminosity because of heavy line blanketing in the v filter. The effective temperature estimate, on the other hand, usually needs only a small correction (e.g. Hubrig, North, & Mathys 2000). The reliability of the gravity estimate from Strömgren photometry is discussed by North & Kroll (1989). From Geneva photometry¹ (Mermilliod, Mermilliod, & Hauck 1997) and using different calibrations (Cramer & Maeder 1979; North & Nicolet 1990; Hauck & North 1993) we obtained effective temperature estimates from 9300 K to 9900 K. These, together with the estimate from *uvby* β , give an average photometrically-determined value of $T_{\text{eff}} = 9600 \pm 300$ K. Interstellar reddening is negligible for this star. There is no clear evidence of any interstellar component in the Na I D1 and D2 lines, which show doublet splitting and belong to the star; any interstellar component is below 0.05 Å in equivalent width. This yields an upper limit to the colour excess of only $E_{B-V} < 0.02$ (Munari & Zwitter 1997).

The Balmer lines profiles are sensitive to effective temperature and gravity. Reliable determination of the right continuum for them – especially for echelle spectra – is not an easy task and may be a source of scatter and errors.

In the case of HD 75049 magnetic broadening is large and needs to be taken into account in the Balmer lines, too. That small variations of Balmer lines may occur has long been known, as discussed recently by, e.g., Valyavin et al. (2007) and Elkin et al. (2008). Balmer lines profiles of H α , H β and H γ in the FEROS spectrum, H β and H γ in the FORS 1 spectra, and H α in the UVES spectra were compared with synthetic profiles for best fits as a function of T_{eff} and $\log g$. An average gives $T_{\text{eff}} = 9700 \pm 170$ K and $\log g = 4.07 \pm 0.28$. The comparison between the observed and synthetic profiles of the Balmer lines gives acceptable agreement for a model with $T_{\text{eff}} = 9600$ K and $\log g = 4.0$. Synthetic calculations of the Balmer profiles were done with the SYNTH code of Piskunov (1992). Model atmospheres by Kurucz (1979), and from the NEMO database (Heiter et al. 2002) were used.

5 MAGNETIC FIELD

5.1 Longitudinal field

From observations with FORS 1 the variable mean longitudinal magnetic field B_z was determined. Table 2 gives the results obtained for three methods of calculation when various lists of spectral lines and spectral regions are used. There are no differences between B_z obtained from all spectral lines in two spectral regions, 3212 – 6215 Å and the shorter 3705 – 6215 Å. But there are shifts of around 600 G between the latter and measurements from the hydrogen lines. These differences most likely reflect the uncertainties involved in the interpretation of the observed circular polarization signal in terms of a magnetic field, which arise from the statistical nature of the measurement (the considered spectral ranges include many lines of different elements, which have different magnetic sensitivities) or from the complexity of the physical foundations of the treatment of the formation of hydrogen lines in A-type star atmospheres in the presence of a magnetic field (Mathys et al. 2000). Differences in the non-uniform distribution of chemical elements in the stellar atmosphere may also contribute, but seem unlikely to represent the dominant effect. The phase curves for all three methods are similar and show sine curves. The distribution

¹ <http://obswww.unige.ch/gcpd/gcpd.html>




Figure 1. Frequency analysis of 13 FORS1 measurements of B_z with times in BJD. The top panel shows the amplitude spectrum with the highest peak at $f = 0.2477 \text{ d}^{-1}$. The middle panel shows the residuals on the same scale after the highest peak has been refined by linear and nonlinear least squares fitting, and prewhitened from the data. It is clear that all of the apparent “noise” in the top panel is spectral window pattern. The bottom panel shows a least-squares fit of a sinusoid to the same data where the best-fitting frequency, $f = 0.2470 \text{ d}^{-1}$, is clear.

of hydrogen in the stellar atmosphere is more homogeneous than for other chemical species. Therefore we used results from hydrogen (last column in Table 2) for further analysis.




Figure 2. Top panel: The variation of the longitudinal magnetic field B_z with rotational period for HD 75049. Filled circles are observations with FORS 1 using the hydrogen lines. The curve is a centred dipole model for $B_p = 42 \text{ kG}$, $i = 25^\circ$ and $\beta = 60^\circ$. Bottom panel: The variation of the magnetic field modulus $\langle B \rangle$ with rotational period for HD 75049 from our UVES and FEROS data. Different symbols are used to represent measurements performed using lines of different ions: filled circles for Fe II $\lambda\lambda 5018 \text{ \AA}$, open squares for Nd III using four lines at $\lambda\lambda 5050 \text{ \AA}$, $5845 \text{ \AA}$, $6145 \text{ \AA}$ and $6327 \text{ \AA}$, and diamonds for Eu II using two lines at $\lambda\lambda 6049 \text{ \AA}$ and $6437 \text{ \AA}$. The solid line is the same model as for upper panel. The dashed curve is a least-squares fit of a sinusoid plus first harmonic to the Fe II data.

5.2 Mean magnetic field modulus

The mean magnetic field modulus $\langle B \rangle$ was determined from high resolution spectra using resolved Zeeman components of several spectral lines. Using Gaussian fitting we determined the centres of shifted Zeeman σ components. The distance between these two components in a spectral line is proportional to value of magnetic field modulus $\langle B \rangle$ (Mathys et al. 1997):

$$\Delta\lambda = 9.34 \times 10^{-13} g_{\text{eff}} \langle B \rangle \lambda_0^2 \quad (3)$$

where wavelength is measured in $\text{\AA}$ and g_{eff} is an effective Landé factor. This relation is valid in fairly general con-

Table 3. The magnetic field modulus $\langle B \rangle$ (kG) in HD 75049 determined from resolved Zeeman components in some selected spectral lines. Because of the non-uniform surface abundance distributions of various ions, some of the differences in field strength at specific rotation phases may be attributable to the distribution of elements. Less reliable results are identified by question marks.

BJD	phase	Fe II 5018 Å	Nd III 5050 Å	Cr II 5237 Å	Nd III 6145 Å	Nd III 6327 Å	Eu II 6437 Å
2454141.66963	0.1550	26.08	26.58	26.83	26.67	26.34	29.49
2454171.50422	0.5234	29.54	28.76	28.23	29.68	28.44	30.10
2454387.81593	0.9470	25.66	26.58	25.91	28.32	26.05	28.78
2454432.82489	0.0631	24.65	25.19?	25.14	25.72	26.08	29.05
2454451.83366	0.7578	27.99	27.98	27.68	29.12	28.01	29.62
2454455.70226	0.7133	28.46	28.29	27.91	29.12	28.38	29.87
2454468.84589	0.9594	25.24	25.82	25.64	28.10	26.08	28.67
2454481.81465	0.1624	26.83	26.64	27.97	27.13	28.73	29.64
2454513.78892	0.0593	24.25	24.74?	24.84?	23.27?		
2454516.68904	0.7755	27.73	27.69	27.48	29.20	27.60	29.52
2454524.66461	0.7453	28.04	27.98	27.83	29.26	28.21	29.62
2454544.51608	0.6481	28.96	28.60	28.18	29.40	28.38	30.06
2454547.51127	0.3878	29.51	28.44	28.57	29.63	29.10	30.04
2454557.59095	0.8773	26.45	26.49	26.71	28.77	26.77	29.98

ditions for lines with doublet or triplet Zeeman patterns; in the general case of more complex, anomalous patterns, it represents only a first approximation. Furthermore, it is based on the assumption that the Doppler effect due to stellar rotation is small compared to the magnetic splitting of the line. The latter is not strictly fulfilled in HD 75049. The resulting asymmetries of the line components, mostly seen between phases 0.85 and 0.35, further hamper the accurate determination of the field modulus.

Since the star is spotted, the actual field strength measured is somewhat dependent on the surface distribution of ions from whose lines the measurements were made. Thus we determined the maximum field strength according to the set of spectral lines used. Table 3 presents our measurements of $\langle B \rangle$ for some spectral lines. Because of the exceptional strength of the magnetic field, the spectral line of Fe II 6149 Å, which lends itself well to the accurate determination of the mean field modulus in Ap stars and has been frequently used to this effect, is strongly distorted by the partial Paschen-Back effect, and its splitting into several components cannot be readily interpreted, so that we did not consider it. In such a strong field, the majority of spectral lines show resolved Zeeman splitting, but many are not strong enough for precise measurements. Many other lines are blends and cannot be used either. The most precise and reliable line splitting measurements were obtained for the strong line Fe II λ 5018 Å. Their mean uncertainties can be estimated from the standard deviation of the individual measurements about a least-squares fit of the superposition of a sinusoid and of its first harmonic against rotation phase. They are found in this way to be of the order of 0.27 kG.

5.3 Dipole model

From Harmanec (1988) we estimate for a main sequence A star with $T_{\text{eff}} = 9600$ K that the stellar radius will be around $2.1 R_{\odot}$. This value agrees well with other published estimations of stellar radii. While main sequence stars with a similar temperature may range in radius between $1.7 R_{\odot}$ and $5 R_{\odot}$ from the zero-age main sequence to the terminal-age

main sequence, a radius more than $2.4 R_{\odot}$ is ruled out for HD 75049, as it requires an implausibly high polar magnetic field in comparison with the observed magnetic field modulus for the dipole model discussed in the next section. The range of possible radius for HD 75049 is therefore between $1.7 R_{\odot}$ and $2.4 R_{\odot}$.

The spectral lines with zero Landé factor, and resolved Zeeman components of lines with large Landé factors, give $v \sin i = 8.5 \pm 1.0 \text{ km s}^{-1}$. From the relation: $v_{\text{eq}} = \frac{50.59 R}{P}$, where P is measured in days, we derive a rotational inclination of $i = 23.6^{\circ} \pm 3.0^{\circ}$ and an equatorial velocity $v_{\text{eq}} = 21.2 \text{ km s}^{-1}$ for $1.7 R_{\odot}$, and $i = 16.5^{\circ} \pm 2.0^{\circ}$ and $v_{\text{eq}} = 30.0 \text{ km s}^{-1}$ for $2.4 R_{\odot}$.

From least squares fitting with the Period04 program (Lenz & Breger 2005), for the hydrogen lines we have an average $B_z = -5.76 \text{ kG}$ which varies with an amplitude of $4.89 \pm 0.06 \text{ kG}$. From expressions for an oblique rotator (e.g., Preston 1967) we find a magnetic obliquity in the range $\beta = 63^{\circ} \pm 3.0^{\circ}$ for star with radius $1.7 R_{\odot}$ to $\beta = 71^{\circ} \pm 3.0^{\circ}$ for $2.4 R_{\odot}$. Following the approach of (Stibbs 1950) and (Preston 1970) we calculated a grid of models using the above parameter range and found a best fit for $B_p = 42 \pm 2 \text{ kG}$, $i = 25^{\circ} \pm 3^{\circ}$ and $\beta = 60^{\circ} \pm 3^{\circ}$, values that are suitable for a stellar radius of $1.7 R_{\odot}$. The variation of the longitudinal magnetic field B_z with the rotational period and selected model fitting is shown in the top panel of Fig. 2. We can get an equally good fit to the top panel of Fig. 2 with the parameters, $R = 2.4 R_{\odot}$, $i = 16.5^{\circ}$, $\beta = 71^{\circ}$ and $B_p = 60 \text{ kG}$, but then the model curve for $\langle B \rangle$ in the bottom panel has a minimum of 38 kG, which is clearly wrong. Under the assumption that the field geometry is dipolar, the implication of this is that HD 75049 is close to the zero-age main sequence.

Variations of $\langle B \rangle$ show less symmetric curves than that for B_z . To fit the $\langle B \rangle$ curve for Fe II 5018 Å we had to employ a sine curve with its first harmonic. Our centred dipole model does not fit the variation of this Fe II line but is acceptable for lines of Nd III and Eu II, considering the errors of measurement and nonuniform distribution of chemical elements. Otherwise the magnetic field geometry is more com-




Figure 3. Variation with rotation phase of the equivalent width (left) and the radial velocity (right) of the lines Cr II λ 5334 Å, Fe II λ 5432 Å, and Nd III λ 6145 Å.

plicated than a simple dipole, as is typical for many magnetic Ap stars (e.g., Landstreet & Mathys 2000) .

5.4 Correlation between magnetic and spectral variability

One sees in Fig. 2 that the field modulus measurements obtained from the three sets of analysed lines show different amplitudes of variation. The Fe-based measurements vary with the largest amplitude, while the lowest variation amplitude is observed for $\langle B \rangle$ values derived from consideration of the Eu lines. These differences may find their origin in part in differences in the non-uniform distribution of the various elements of interest over the stellar surface. In order to test this interpretation, we have checked the dependence on rotation phase of the equivalent widths and radial velocities of lines of the considered elements. The examples in Fig. 3 illustrate the observed behaviours. The spectral lines that are shown, Cr II λ 5334 Å, Fe II λ 5432 Å and Nd III λ 6145 Å, are representative of all the reasonably unblended lines of the elements from which they arise. Equivalent width variations are definitely observed for Cr and Nd, but they are not definitely detected for Fe. The Nd lines also show radial velocity

variations, while our data show no clear indications of such variations for Cr and Fe. Even after averaging the radial velocity measurements of several lines of these elements, no convincing evidence for variability was found.

For Nd, maximum equivalent width occurs close to phase 0.5, hence to the negative extremum of the longitudinal field, and to the maximum of the field modulus. The radial velocities of the Nd lines vary in phase quadrature with their equivalent widths; the lines are blueshifted between phases 0 and 0.5, and redshifted between phases 0.5 and 1.0. This is consistent with the presence of a region of enhanced abundance (“spot”) of Nd around the negative magnetic pole. The same conclusion applies to Eu, whose lines show similar variability to those of Nd.

The equivalent widths of the Cr lines show the opposite variation, reaching their minimum close to phase 0.5, and being largest close to phase 0. Any radial velocity variations that they may show do not clearly stand out of the measurement noise, so that their amplitude is definitely much lower (by a factor of 5 or more) than for Nd lines. Yet, this may not be inconsistent with equivalent width variations being due to a non-uniform distribution of Cr over the stellar surface such that the abundance of this element is minimum around the

negative pole, and increases away from it. Possible configurations include a band of enhanced Cr abundance along the magnetic equator. With a distribution of this type, one can simultaneously observe on the visible stellar hemisphere approaching and receding Cr-rich regions, whose contributions to the radial velocity of Cr spectral lines mostly averages out in disk-integrated observations.

Finally, Fe lines do not show any definite variability of their equivalent widths or radial velocities, indicating that any abundance inhomogeneity in the distribution of this element on the visible part of the star must be moderate at most. Si behaves like Fe.

Thus we are left with the qualitative picture of a star showing a concentration of Nd and Eu and a (relative) depletion of Cr around its negative magnetic pole, with a comparatively uniform distribution of Fe and Si over its surface. One should keep in mind, though, that due to the low inclination of the rotation axis on the line of sight, a large fraction of the stellar surface is never observed.

The non-uniform distribution of Nd over the stellar surface may account for part of the differences between the values of $\langle B \rangle$ that are derived from measurements of its lines, and from those of Fe, whose abundance appears much more constant across the star. However, it is unlikely to be the only factor, or even the main one. If it were, one would, for instance, expect greater values of the field modulus to be determined from Nd lines than from Fe lines close to phase 0.5; the opposite is observed in Fig. 2. Most likely, the origin of the difference should be sought elsewhere. In particular, one should bear in mind that Nd lines are subject to hyperfine structure. This introduces severe complications in the treatment of their formation in the presence of a strong magnetic field, as illustrated, e.g., in Landi Degl’Innocenti (1975). Accordingly, the physical meaning of the value of $\langle B \rangle$ that is obtained by application of Eq. (3) to the observed splitting of Nd lines is not fully clear. In particular, one cannot a priori expect it to provide a measurement of the mean magnetic field modulus that is consistent with its determinations from analysis of Fe lines, which rests on a firmer physical basis. The same applies to the usage of Eu lines to measure $\langle B \rangle$. The differences seen in Fig. 2 between the variation curves of this field moment resulting from measurements of the Nd III and Eu II lines would actually be very difficult to understand in terms of abundance inhomogeneities since the equivalent width and radial velocity behaviours indicate that, to first order, the distribution of both elements over the stellar surface is similar. On the other hand, around phase 0, the lines Eu II $\lambda\lambda 6049 \text{ \AA}$ and $6437 \text{ \AA}$ become very weak and shallow, so that the uncertainties of the $\langle B \rangle$ determinations based on them become very large, and the values obtained for this field moment between phases ~ 0.9 and ~ 0.2 should be considered with appropriate caution.

6 ASYMMETRY OF SPECTRAL LINES

Spectral lines in HD 75049 show significant variability with rotation period, and for many lines we also note asymmetry in the Zeeman patterns, as can be seen in Fig. 4. Shifted σ components normally are symmetric with respect to the line centre, according to classical Zeeman theory. Observations of magnetic Ap stars reveal more complicated behaviours

of the Zeeman patterns. Spectral line asymmetry is often visible in high resolution spectra of Ap stars. While non-uniform distribution of chemical species on the stellar surface contributes to this, the main source of line asymmetry more often is the variable combination of Zeeman and Doppler effects across the stellar surface. Indeed, due to the large-scale structure of the magnetic field, the Zeeman and Doppler shifts are correlated across the stellar disk, and this correlation is reflected in the shapes of the disk-integrated spectral lines.

The observed presence of this effect in HD 75049 is particularly plausible, considering that this star has a very strong magnetic field, and a large $v \sin i$ for a star with magnetically resolved lines. That it is the dominant effect for explanation of the origin of the line asymmetries can be inferred from consideration of Fig. 4. Representative lines of each of the three elements Fe, Cr, and Nd are seen to show qualitatively similar asymmetries: a red σ component (or red wing) shallower and broader than its blue counterpart at phases comprised between ~ 0.6 and ~ 0.9 , and vice-versa in the phase range $\sim 0.1 - 0.4$. Since the distributions of Fe, Cr and Nd over the star differ considerably from each other, the similarity of the line asymmetries for all of them cannot arise primarily from their inhomogeneities. Also, because the magnetic field of HD 75049 is very strong, partial Paschen-Back effect may contribute to asymmetries of a significant number of lines (Mathys 1990), but it affects different transitions to a variable extent and in qualitatively different ways, so that it cannot account for the consistent behaviours illustrated in Fig. 4.

7 CHEMICAL ABUNDANCES

Large overabundances of some chemical elements, especially of rare earth elements, is one of the defining characteristics of Ap stars. Various chemical elements concentrate in different places on the surface of the star. There is a tendency for rare earth elements to concentrate around the magnetic poles in some sort of spots. The size and number of spots may vary (Kochukhov et al. 2004; Freyhammer et al. 2009). The situation is even more complex since the atmospheres of Ap stars also show vertical stratification (e.g., Babel & Lanz 1992; Wade et al. 2001; Kochukhov et al. 2009) where various elements accumulate in different layers in the atmosphere.

Because of the spotted structure of Ap stars, average abundances do not reflect the complexity of the physical conditions at the stellar surface, yet they are useful for statistical analysis and comparison of stars in the class (e.g., Adelman 1973; Ryabchikova et al. 2004). HD 75049 has a rich spectrum, but is not extraordinarily peculiar. In the spectrum many lines of rare earth elements are visible, but the majority of the strongest metal lines belong to Si II, Ti II, Fe II and Cr II. Lines of Sr II in the blue region of the FEROS spectrum are also rather strong.

We obtained the abundances of several elements using spectral synthesis where the observed spectra were compared with synthetic spectra until a best fit was obtained. An example for Fe II $\lambda 5018 \text{ \AA}$ is shown in Fig. 5. The synthetic spectra were calculated with the software SYNTHMAG (Piskunov 1999). The spectral line list was taken from




Figure 4. Variation with rotation phase of the observed profiles of the spectral lines Fe II λ 5018 Å, Nd III λ 5050 Å, and Cr II λ 5334 Å. Each spectrum is shifted vertically from the normalised continuum to a value corresponding to $(1 + \text{rotation phase})$, hence the ordinate shows both continuum and rotation phase.

Table 4. Chemical abundances for HD 75049 for selected elements, and their corresponding solar abundances (Asplund, Grevesse, & Sauval 2005). The errors quoted are internal standard deviations for the set of lines measured. Only an upper limit could be placed on the abundance of Nd II. Columns 3 and 5 give the number of lines used, N_{lines} , in each case.

Ion	$\log N/N_{\text{tot}}$ $\phi = 0.388$	N_{lines}	$\log N/N_{\text{tot}}$ $\phi = 0.063$	N_{lines}	$\log N/N_{\text{tot}}$ Sun
Si II	-3.70 ± 0.05	2	-3.55 ± 0.05	2	-4.49
Ti II	-6.04 ± 0.10	2	-5.85 ± 0.15	2	-7.10
Cr II	-4.73 ± 0.15	14	-4.53 ± 0.13	9	-6.36
Fe I	-3.93 ± 0.17	10	-4.10 ± 0.29	4	-4.55
Fe II	-3.89 ± 0.15	17	-3.98 ± 0.15	6	-4.55
La II	-7.15 ± 0.21	16	-7.23 ± 0.18	6	-10.87
Ce II	-7.48 ± 0.16	5	-7.95 ± 0.05	4	-10.42
Pr III	-8.07 ± 0.15	4	-8.10 ± 0.16	4	-11.29
Nd III	-7.27 ± 0.05	3	-7.57 ± 0.12	3	-10.55
Nd II	< -8.40	3	< -7.98	3	-10.55
Eu II	-7.71 ± 0.19	4	-8.10 ± 0.29	4	-11.48

the Vienna Atomic Line Database (VALD, Kupka et al. 1999), which includes lines of rare earth elements from the DREAM database (Biémont, Palmeri, & Quinet 1999). A model atmosphere with $T_{\text{eff}} = 9600$ K and $\log g = 4.0$ with metallicity of 0.5 dex above solar from the NEMO database (Heiter et al. 2002) was used. The synthetic spectra provided a good match to many spectral lines with the field strength chosen, but some lines need further broadening to match the wings. Macroturbulence is possibly not the explanation for this, as magnetic fields suppress macroscopic motion in the stellar atmospheres. With the non-uniform distribution of abundances – both horizontally and vertically – in Ap star atmospheres, it may be that stronger magnetic field strengths in different line-forming regions could explain

this extra broadening. Further study is needed to test this idea.

Two spectra at rotation phases near minimum and maximum longitudinal field strength (phase 0.063 and 0.388; see Fig. 2) were used for abundances determination. Magnetic field strengths of 26 kG and 30 kG, respectively, were used for calculation of the synthetic spectra at these two rotational phases. Synthetic line profiles for various abundances were compared with observed profiles for best fits. Many calculated line profiles do not fit the observed profiles perfectly as a result of blending and asymmetry, but the average value for a number of lines gives reliable results. It was more difficult to fit the spectrum observed at rotational phase 0.063 when the field strength is lower and some lines are not as sharp and more asymmetric. Table 4 presents the




Figure 5. Observed (solid line) and synthetic (dashed line) profiles for Fe II 5018 Å.

values of the abundances with errors determined for various ions. The errors given are the internal standard deviations for the list of examined lines for each ion. There were no lines of Nd II found in the spectra, so only an upper limit is given in Table 4. The last column in the table gives solar abundances for the same elements taken from Asplund, Grevesse, & Sauval (2005). HD 75049 shows overabundances of silicon, chromium and europium and thus can be considered to belong to the SiCr group of Ap stars, according to the classification of peculiar stars.

8 CONCLUDING REMARKS

The strong magnetic field of HD 75049 is an exciting discovery. This star is a rival for the long-known strongest Ap magnetic field in Babcock's star, HD 215441. Both stars have deep silicon lines, but are at opposite ends of the temperatures range of the Ap Si stellar group. Babcock (1960) noted that HD 215441 does not have outstanding peculiarities in the spectrum. We have found the same in HD 75049, which has a rich peculiar spectrum, but with overabundances of number chemical elements similar to other Ap stars with much smaller field strengths.

HD 215441 has a magnetic field modulus that changes from 32 to 35 kG over the rotational period (Babcock 1960; Preston 1969). Preston (1969) suggested that the magnetic field of this star deviates from a centred dipole. Magnetic configurations with a combination of dipole and quadrupole components (Borra & Landstreet 1978) or dipole, quadrupole and octupole (Landstreet & Mathys 2000) components were required to model both observations of B_z obtained by Borra & Landstreet (1978) and $\langle B \rangle$ by Preston (1969).

In contrast, for HD 75049 the centred dipole model is a suitable first approximation to fit both the longitudinal field

and field modulus measurements. Some discrepancy with the second magnetic moment dipole model may be a result of nonuniform distribution of chemical elements at surface. More complex magnetic geometry may also be present in this star. An important result is that the star is probably young. Its estimated stellar radius places it close to the zero-age main sequence.

Landstreet & Mathys (2000) examined the inclination of rotation axis i to line-of-sight and obliquity β of the magnetic axis to the rotation axis for a sample of magnetic Ap stars. They found that the majority of the slow rotators with periods longer than 25 d have β smaller than 20° , while stars with short rotation periods tend to show a large obliquity of the magnetic axis. Our estimation of $\beta = 60^\circ \pm 3^\circ$ for HD 75049 is consistent with this conclusion.

9 ACKNOWLEDGEMENTS

We thank Dr. Romanyuk for providing information about the Zeeman configuration of some spectral lines and for useful discussion. DWK and VGE acknowledge support for this work from the Science and Technology Facilities Council (STFC) of the UK. This research has made use of NASA's Astrophysics Data System and SIMBAD database, operated at CDS, Strasbourg, France.

REFERENCES

- Abt H. A., 2009, AJ, 138, 28
- Adelman S. J., 1973, ApJ, 183, 95
- Asplund M., Grevesse N., Sauval A. J., 2005, ASPC, 336, 25
- Babcock H. W., 1947, ApJ, 105, 105
- Babcock H. W., 1958, ApJS, 3, 141
- Babcock H. W., 1960, ApJ, 132, 521
- Babel J., Lanz T., 1992, A&A, 263, 232
- Biémont E., Palmeri P., Quinet P., 1999, Ap&SS, 269, 635
- Borra E. F., Landstreet J. D., 1978, ApJ, 222, 226
- Cramer N., Maeder A., 1979, A&A, 78, 305
- Elkin V. G., Kurtz D. W., Freyhammer L. M., Hubrig S., Mathys G., 2008, MNRAS, 390, 1250
- Freyhammer L. M., Elkin V. G., Kurtz D. W., Mathys G., Martinez P., 2008, MNRAS, 389, 441
- Freyhammer L. M., Kurtz D. W., Elkin V. G., Mathys G., Savanov I., Zima W., Shibahashi H., Sekiguchi K., 2009, MNRAS, 396, 325
- Harmanec P., 1988, BAICz, 39, 329
- Hauck B., North P., 1993, A&A, 269, 403
- Heiter U., et al., 2002, A&A, 392, 619
- Hubrig S., et al., 2005, A&A, 440, L37
- Hubrig S., Mathys G., Kurtz D. W., Schöller M., Elkin V. G., Henrichs H. F., 2009, MNRAS, 657
- Hubrig S., North P., Mathys G., 2000, ApJ, 539, 352
- Hubrig S., North P., Schöller M., Mathys G., 2006, AN, 327, 289
- Kochukhov O., 2006, A&A, 454, 321
- Kochukhov O., Drake N. A., Piskunov N., de la Reza R., 2004, A&A, 424, 935
- Kochukhov, O., Shulyak, D., & Ryabchikova, T. 2009, A&A, 499, 851

- Kudryavtsev D. O., Romanyuk I. I., Elkin V. G., Paunzen E., 2006, MNRAS, 372, 1804
- Kupka F., Piskunov N., Ryabchikova T. A., Stempels H. C., Weiss W. W., 1999, A&AS, 138, 119
- Kurtz D. W., 1985, MNRAS, 213, 773
- Kurtz D. W., Elkin V. G., Cunha M. S., Mathys G., Hubrig S., Wolff B., Savanov I., 2006, MNRAS, 372, 286
- Kurucz R. L., 1979, ApJS, 40, 1
- Landi Degl'Innocenti, E., 1975, A&A, 45, 269
- Landstreet J. D., Mathys G., 2000, A&A, 359, 213
- Lenz P., Breger M., 2005, CoAst, 146, 53
- Martinez P., 1993, PhD Thesis, University of Cape Town
- Mathys G., 1990, A&A, 232, 151
- Mathys G., 1991, A&AS, 89, 121
- Mathys G., 1995, A&A, 293, 746
- Mathys G., Hubrig S., Landstreet J. D., Lanz T., Manfroid J., 1997, A&AS, 123, 353
- Mathys G., Stehlé C., Brillant S., Lanz T. 2000, A&A, 358, 1151
- Mermilliod J.-C., Mermilliod M., Hauck B., 1997, A&AS, 124, 349
- Moon T. T., Dworetzky M. M., 1985, MNRAS, 217, 305
- Munari U., Zwitter T., 1997, A&A, 318, 269
- Napiwotzki, R., Schoenberner, D., & Wenske, V. 1993, A&A, 268, 653
- North P., Kroll R., 1989, A&AS, 78, 325
- North P., Nicolet B., 1990, A&A, 228, 78
- Pojmanski G., 2002, AcA, 52, 397
- Piskunov, N. E. 1992, Stellar Magnetism, 92
- Piskunov N., 1999, ASSL, 243, 515
- Preston G. W., 1967, ApJ, 150, 547
- Preston G. W., 1969, ApJ, 156, 967
- Preston G. W., 1970, ApJ, 160, 1059
- Ryabchikova T., Nesvacil N., Weiss W. W., Kochukhov O., Stütz C., 2004, A&A, 423, 705
- 1992,
- Stibbs D. W. N., 1950, MNRAS, 110, 395
- Valyavin G., Lee B.-C., Shulyak D., Han I., Kochukhov O., Khang D.-I., Kim K.-M., 2007, ASPC, 362, 245
- Wade G. A., Ryabchikova T. A., Bagnulo S., Piskunov N., 2001, ASPC, 248, 373